

DENOSUMAB (Xgeva) - local funding required

INDICATION (ICD10) D48.1

1. Neoadjuvant and maintenance single agent in giant cell tumours of the bone

REGIMEN

Cycle 1

Days 1, 8 and 15 DENOSUMAB 120mg SC once daily

Cycle 2 onwards

Day 1 DENOSUMAB 120mg SC once daily

CYCLE FREQUENCY AND NUMBER OF CYCLES

Every 28 days

ANTI-EMETICS

None required

CONCURRENT MEDICATION REQUIRED

Denosumab	Vitamin D. Adequate intake of calcium
-----------	--

INVESTIGATIONS

Blood results required before SACT administration

FBC, U&E, Ca⁺⁺ and LFTs every cycle

Bone profile including Vitamin D, PTH) PINP every cycle

Neutrophils x 10⁹/L ≥1.5

Platelets x 10⁹/L ≥100

Serum creatinine every cycle

Baseline weight and every cycle

MAIN TOXICITIES AND ADVERSE REACTIONS

Denosumab	Osteonecrosis of jaw/auditory canal, suspend treatment if invasive dental work needed. Hypocalcaemia must be corrected by adequate intake of calcium and vitamin D before initiating therapy. Patients with severe renal impairment (CrCl <30ml/min) or receiving dialysis are at greater risk of developing hypocalcaemia. Infections and infestations - UTI, respiratory, diverticulitis, cellulitis and ear infections Sciatica Cataracts Constipation Skin - rash and eczema Pain
-----------	--

INTERACTIONS WHICH MAY REQUIRE DOSE MODIFICATIONS

(not exhaustive list check SPC/BNF/Stockleys)

Denosumab	-
-----------	---



DOSE MODIFICATIONS

Hepatic impairment

Not studied

Renal impairment

No dose adjustment required

REFERENCES

1. Thomas D et al Denosumab in patients with giant-cell tumour of bone; an open label phase 2 study. Lancet Oncology vol11 March 2010; 275-280