

## ATEZOLIZUMAB (Tecentriq) CARBOPLATIN ETOPOSIDE

### INDICATION (ICD10) C34

Check the most recent *Blueteq* eligibility criteria before prescribing. *Blueteq* registration required. ([www.england.nhs.uk/publication/national-cancer-drugs-fund-list/](http://www.england.nhs.uk/publication/national-cancer-drugs-fund-list/)) (ATE7)

1. For the first-line treatment of adult patients with extensive-stage small cell lung cancer (Not received previous systemic therapy for his/her extensive stage disease. Previous treatment with concurrent chemoradiotherapy for limited stage SCLC is allowed as long as therapy was completed at least 6 months prior to the diagnosis of recurrent and extensive stage disease) No symptomatically active brain metastases or leptomeningeal metastases. PS 0 or 1. (TA638)

### REGIMEN SC

Cycles 1 to 4

Day 1	ATEZOLIZUMAB	1875mg SC
	CARBOPLATIN	AUC 5 in #ml glucose 5% IV infusion over 30 minutes Dose calculated by EDTA GFR or calculated CrCl + 25 x AUC. (Maximum dose when using CrCl 125+25 x AUC)
	ETOPOSIDE	100mg/m <sup>2</sup> in #ml sodium chloride 0.9% IV infusion over 60 minutes
Day 2	ETOPOSIDE	100mg/m <sup>2</sup> in #ml sodium chloride 0.9% IV infusion over 60 minutes
Day 3	ETOPOSIDE	100mg/m <sup>2</sup> in #ml sodium chloride 0.9% IV infusion over 60 minutes

### Cycles 5 onwards

Day 1 ATEZOLIZUMAB 1875mg SC

# diluent volume for dose prescribed as per national standardised product specification

Atezolizumab 1875mg SC – Administer in the thigh over 7 minutes. Alternate thighs, administer at least 2.5cm from the old site.

### REGIMEN IV

Cycles 1 to 4

Day 1	ATEZOLIZUMAB	1200mg in #ml sodium chloride 0.9% IV infusion
	CARBOPLATIN	AUC 5 in #ml glucose 5% IV infusion over 30 minutes Dose calculated by EDTA GFR or calculated CrCl + 25 x AUC. (Maximum dose when using CrCl 125+25 x AUC)
	ETOPOSIDE	100mg/m <sup>2</sup> in #ml sodium chloride 0.9% IV infusion over 60 minutes
Day 2	ETOPOSIDE	100mg/m <sup>2</sup> in #ml sodium chloride 0.9% IV infusion over 60 minutes
Day 3	ETOPOSIDE	100mg/m <sup>2</sup> in #ml sodium chloride 0.9% IV infusion over 60 minutes

### Cycles 5 onwards

Day 1 ATEZOLIZUMAB 1200mg in #ml sodium chloride 0.9% IV infusion

# diluent volume for dose prescribed as per national standardised product specification

Atezolizumab 1200mg IV – The initial dose should be delivered over 60 minutes.

If the first infusion is tolerated without infusion-associated adverse events, the second infusion may be delivered over 30 minutes.

If the 30-minute infusion is well tolerated, all subsequent infusions may be delivered over 30 minutes.

Etoposide - days 2 and 3 can be given orally ETOPOSIDE 200mg/m<sup>2</sup>/day but is not recommended as oral absorption is variable (it may cause reduced efficacy or severe toxicity in patients), the intravenous route is preferred. However for logistical reasons the oral route may be necessary.

## CYCLE FREQUENCY AND NUMBER OF CYCLES

Combination every 21 days for 4 cycles

Atezolizumab monotherapy from cycle 5 may continue until disease progression.

## ANTI-EMETICS

Moderate emetic risk day 1 cycles 1 to 4

Low emetic risk days 2 and 3 cycles 1 to 4

Low risk day 1 cycle 5 onwards

## CONCURRENT MEDICATION REQUIRED

Carboplatin	Anaphylaxis treatment should be prescribed if the patient has had an anaphylactic episode previously. Dexamethasone 20mg IV bolus Chlorphenamine 10mg IV bolus Carboplatin should be given at a slower rate e.g. 2-4 hours.
-------------	--

## EXTRAVASATION AND TYPE OF LINE / FILTERS

Atezolizumab IV – neutral

Carboplatin – irritant

Etoposide - irritant

Atezolizumab IV use of 0.2-5micron filter is optional

Peripheral line

## INVESTIGATIONS

Blood results required before SACT administration

FBC, U&E and LFTs every cycle

Neutrophils x 10<sup>9</sup>/L ≥1.5

Platelets x 10<sup>9</sup>/L ≥100

Ideally EDTA GFR should be used

Creatinine clearance (GFR) calculated, at the Consultants discretion

Serum creatinine - each cycle

Thyroid function baseline and every 1 to 2 cycles

Random cortisol baseline and every 1 to 2 cycles

Baseline weight and every cycle

## MAIN TOXICITIES AND ADVERSE REACTIONS

Atezolizumab	Immune mediated pneumonitis Immune mediated hepatitis Immune mediated colitis Immune mediated endocrinopathies
Carboplatin	Ototoxicity - monitor Neurotoxicity – monitor.

## INTERACTIONS WHICH MAY REQUIRE DOSE MODIFICATIONS

(not exhaustive list check SPC/BNF/Stockleys)

Carboplatin	Aminoglycosides increased risk of nephrotoxicity and ototoxicity. Renal function should be well monitored and audiometric tests as required. Carboplatin can cause a decrease in phenytoin serum levels. This may lead to reappearance of seizures and may require an increase of phenytoin dosages.
-------------	---

## DOSE MODIFICATIONS

### Non-haematological

#### Atezolizumab

Immune-related adverse reactions - refer to TV immune-oncology agent immune related adverse event clinical guideline.

If the drug-related toxicity does not resolve to grade 0-1 within 12 weeks after onset of toxicity, discontinuation is recommended.

### Hepatic impairment

#### Etoposide

Bilirubin 26-51micromol/L or AST 60-180u/L	give 50% dose
Bilirubin >51micromol/L or AST >180u/L	Clinical decision

### Renal impairment

#### Carboplatin

GFR / calculated CrCl $\leq$ 20ml/min or $\leq$ 30ml/min with pre-existing severe renal impairment	contraindicated
--	-----------------

#### Etoposide

CrCl >50ml/min	give 100% dose
CrCl 15-50ml/min	give 75% dose
CrCl <15ml/min	further dose reduction required

## REFERENCES

1. Horn, L et al; NEJM 2018; 379: 2220 - 2229