

BEVACIZUMAB IRINOTECAN OXALIPLATIN Modified de Gramont (FOLFOXIRI)

INDICATION (ICD10) C18, C20

Check the most recent *Blueteq* eligibility criteria before prescribing. *Blueteq* registration required (www.england.nhs.uk/publication/national-cancer-drugs-fund-list/) (BEV11)

1. Bevacizumab with FIRST LINE fluoropyrimidine-based chemotherapy for metastatic or locally advanced and inoperable colorectal cancer and has not received any previous systemic therapy for this indication. Patients may have received neoadjuvant systemic therapy for non-metastatic disease and/or adjuvant chemotherapy after surgery and has a documented presence of microsatellite stability (MSI-S) or DNA mismatch repair proficiency (pMMR) confirmed by validated testing, OR immunotherapy is not being used as first line therapy due to its unsuitability for this patient. The primary reason for the patient NOT receiving either cetuximab or panitumumab alongside first line chemotherapy is has a right sided primary tumour, the tumour has a mutant RAS status or the RAS test result not yet reported and the decision to proceed without knowing RAS status has been discussed with the patient during the consenting process or cetuximab or panitumumab are not suitable for this patient due to pre-existing medical conditions or sensitivities.

REGIMEN

Day 1	BEVACIZUMAB	5mg/kg	IV infusion	#ml sodium chloride 0.9%
	Premedication 30 minutes prior to irinotecan: Atropine 250mcg subcutaneously			
	IRINOTECAN	165mg/m ²	IV infusion	#ml diluent over 30 minutes
	CALCIUM FOLINATE	350mg	IV infusion	250ml glucose 5% over 2 hours concurrently with oxaliplatin
	OXALIPLATIN	85mg/m ²	IV infusion	#ml glucose 5% over 2 hours
	FLUOROURACIL	3200mg/m ²	IV infusion	continuous over 46 hours

diluent and diluent volume for dose prescribed as per national standardised product specification or licensed dose

NB Calcium folinate (calcium leucovorin) is not the same as calcium levofolinate. Calcium levofolinate is a single isomer of folinic acid and the dose is generally half that of calcium folinate.

CYCLE FREQUENCY AND NUMBER OF CYCLES

Every 14 days until disease progression

ANTI-EMETICS

Highly emetogenic day 1
Low emetogenic risk day 2

CONCURRENT MEDICATION REQUIRED

Bevacizumab	None
Fluorouracil	Mouth and bowel support to be considered eg loperamide, benzydamine mouthwash
Irinotecan	Ensure premedication atropine given 30 minutes prior to treatment
Oxaliplatin	Flush with glucose 5% before and after infusion
GCSF	Consider GCSF

ADMINISTRATION

Bevacizumab	The initial dose should be administered over 90 minutes, if tolerated well the second infusion may be administered over 60 minutes. If the 60 minute infusion is well tolerated all subsequent infusions may be administered over 30 minutes.
-------------	--

EXTRAVASATION AND TYPE OF LINE / FILTERS

Bevacizumab - neutral
Fluorouracil – inflammitant
Irinotecan - irritant
Oxaliplatin – exfoliant

Filter not required
Central line (single lumen)

INVESTIGATIONS

Blood results required before SACT administration

FBC, U&E including Mg ⁺⁺ and LFTs Neutrophils $\geq 1.5 \times 10^9/L$ Platelets $\geq 100 \times 10^9/L$	baseline and every cycle
EDTA GFR or calculated CrCl at consultant's discretion.	baseline and every cycle
Serum creatinine	baseline and every cycle
Blood pressure	baseline and before every bevacizumab dose
Urinalysis for proteinuria	baseline and before every bevacizumab dose
DPYD (dihydropyrimidine dehydrogenase) test	baseline if no previous result / treatment with a fluoropyrimidine
Weight	baseline and every cycle

Formatted Table

MAIN TOXICITES AND ADVERSE REACTIONS

Bevacizumab	Arterial thromboembolism Gastrointestinal perforation Haemorrhage Hypertension Wound healing complications
Fluorouracil	Palmar plantar (handfoot syndrome) causing red palms and soles – treat with pyridoxine 50mg tds. Diarrhoea – treat with loperamide or codeine Cardiotoxicity – monitor cardiac function (consider ECG at baseline). Special attention is advisable in treating patients with a history of heart disease, arrhythmias or angina pectoris or those who develop chest pain during treatment with fluorouracil. Stomatitis
Irinotecan	Acute cholinergic syndrome (including diarrhea and delayed diarrhea, abdominal pain, hypotension, dizziness, malaise, increased salivation). Drink large volumes of fluid containing electrolytes and an appropriate antiarrhoeal therapy - loperamide 4mg initially then 2mg every 2 hours, continuing for 12 hours after the last liquid stool (maximum of 48 hours in



	total).
Oxaliplatin	Peripheral sensory neuropathy and laryngeal spasm – avoid cold drinks and touching cold items

INTERACTIONS WHICH MAY REQUIRE DOSE MODIFICATIONS
(not exhaustive list check SPC/BNF/Stockleys)

Bevacizumab	-
Fluorouracil	Cimetidine slightly increases exposure to fluorouracil Metronidazole increased toxicity Phenytoin concentration increased Warfarin
Irinotecan	Aprepitant and fosaprepitant increases exposure to irinotecan. Carbamazepine and St John's wort decreases exposure to irinotecan, avoid. Enzalutamide, mitotane, phenobarbitone, phenytoin, primidone and rifampicin decreases exposure to irinotecan, avoid.

DOSE MODIFICATIONS

Haematological

Neutrophils $<1.5 \times 10^9/L$ and/or platelet count $<100 \times 10^9/L$	delay one week, only treat when neutrophils and platelets are above these limits.
Grade 4 neutropenia	consider giving 50% irinotecan and fluorouracil in palliative disease or GCSF support for non-palliative disease.
If >1 delay or 1 delay ≥ 2 weeks reduce all the irinotecan, oxaliplatin and fluorouracil doses to give 80% for future cycles. A further dose reduction may be made at the clinician's discretion.	

Non-haematological

Bevacizumab

Hypertension

Baseline blood pressure should be $<150/100$ mmHg.

Diastolic increase >20 mmHg above baseline or BP rises to $>150/100$ mmHg	Antihypertensive therapy may be required.
Blood pressure $>180/110$ mmHg	It is advised that bevacizumab therapy is withheld until blood pressure controlled.

Proteinuria

Urine dipstick result. 1+ or 2+ on dipstick (0.3–2.9g/L)	Continue with bevacizumab. No additional evaluation required.
3+ on dipstick (3–19g/L)	May have dose of bevacizumab as scheduled, but 24 hour urine to measure 24 hour protein to be done a few days before next cycle due. If 24hr protein result <2 g, continue with bevacizumab, with continued proteinuria monitoring via 24 hour urine before each dose. If the 24 hour protein level falls to <1 g/24hr, return to dipstick analysis. If ≥ 2 g, withhold bevacizumab until repeat 24 hour urine collection shows <2 g protein. Then re-introduce bevacizumab, with continued proteinuria monitoring via 24 hour urine.
4+ on dipstick (≥ 20 g/L)	Withhold bevacizumab. 24 hour urine required.

Follow 24 hour urine monitoring and guidance as for 3+ on dipstick.

Wound healing

Bevacizumab may adversely affect the wound healing process. Therapy should not be initiated for at least 28 days following major surgery or until the surgical wound is fully healed. Therapy should also be withheld for at least 28–60 days before elective surgery.

Irinotecan

If patients suffer from severe diarrhoea, which required IV rehydration or neutropenic fever, consider reduction in subsequent cycles, discuss with SpR or Consultant.

Oxaliplatin

If patients develop acute laryngopharyngeal dysaesthesia infuse the next cycle over 4 hours.

If symptoms persist give 80% dose.

If persistent sensory symptoms occur, withdraw treatment

Hepatic impairment

Bevacizumab

The safety and efficacy have not been studied in patients with hepatic impairment.

Fluorouracil

Significantly impaired hepatic function eg bilirubin >50micromol/L may be a sign of disease progression and require cessation of, or change in, treatment. Always discuss deteriorating liver function with consultant.	
If hepatic function is impaired, the recommended dose can be reduced to give 50% to 70% dose, but no need for dose adjustment is expected in mild and moderate (without renal impairment).	
Bilirubin >85micromol/L	not recommended

Irinotecan

Bilirubin 24-50micromol/L	give 50% dose
Bilirubin >51micromol/L	not recommended

Oxaliplatin

No dose adjustment is needed.

Renal impairment

Bevacizumab

The safety and efficacy have not been studied in patients with **hepatic-renal** impairment.

Fluorouracil

If renal function is impaired, the recommended dose can be reduced to give 50% to 70% dose, but no need for dose adjustment is expected.
--

Irinotecan

Not recommended in renal impairment, use with caution.

Oxaliplatin

CrCl >30ml/min	give 100% dose
CrCl <30ml/min	Dose reduce (consider 50% of original dose)



REFERENCES

1. CDF list

Assessments

	Pre	Cycle 1	Cycle 2	Cycle 3	Cycle 4	Ongoing
Clinical assessment	X		Pre cycle		Pre cycle	Alternate cycles or team discretion
SACT assessment (PS and toxicities)	X	X	X	X	X	Every cycle
FBC	X	X	X	X	X	Every cycle
U&E, calcium, magnesium & LFT	X	X	X	X	X	Every cycle
CrCl	X	X	X	X	X	Every cycle
<u>Blood pressure</u>	<u>X</u>	<u>X</u>	<u>X</u>	<u>X</u>	<u>X</u>	<u>Every cycle</u>
<u>Urine protein</u>	<u>X</u>	<u>X</u>	<u>X</u>	<u>X</u>	<u>X</u>	<u>Every cycle</u>
Dihydropyrimidine dehydrogenase (DPYD) deficiency test	X					This test is normally only required if a patient has not had capecitabine or fluorouracil in the past. However a consultant may still request this test if capecitabine or fluorouracil was not tolerated previously. The result MUST be available before administration of chemotherapy unless clear documentation from the consultant is available to the contrary. DPD deficiency may lead to life threatening toxicity.
CT scan	X					By cycle 6, Inform consultant team if not booked
Informed consent	X					Verbal each cycle
Height	X					
Weight recorded	X	X	X	X	X	Every cycle